



ACUTE MYELOID LEUKEMIA: MOLECULAR ONCOLOGY TUMOR BOARDS COURSE REFERENCES

Arber DA, Orazi A, Hasserjian R, Thiele J, Borowitz MJ, Le Beau MM, et al. The 2016 revision to the World Health Organization classification of myeloid neoplasms and acute leukemia. *Blood*. 2016;127(20):2391-405. ([View Abstract](#))

Brunet S, Labopin M, Esteve J et al. Impact of FLT3 internal tandem duplication on the outcome of related and unrelated hematopoietic transplantation for adult acute myeloid leukemia in first remission: a retrospective analysis. *J Clin Oncol*. 2012 Mar 1;30(7):735-41. ([View Abstract](#))

Cancer Genome Atlas Research N. Genomic and epigenomic landscapes of adult de novo acute myeloid leukemia. *N Engl J Med*. 2013;368(22):2059-74. ([View Abstract](#))

Chen W, Konoplev S, Medeiros LJ, Koeppen H, Leventaki V, Vadhan-Raj S, et al. Cuplike nuclei (prominent nuclear invaginations) in acute myeloid leukemia are highly associated with FLT3 internal tandem duplication and NPM1 mutation. *Cancer*. 2009;115(23):5481-9. ([View Abstract](#))

Cooper BW, Kindwall-Keller TL, Craig MD, et al. A phase I study of midostaurin and azacitidine in relapsed and elderly AML patients. *Clin Lymphoma Myeloma Leuk*. 2015 Jul;15(7):428-432. ([View Abstract](#))

Cortes J, Perl A, Dombret H, et al. Final Results of a Phase 2 Open-Label, Monotherapy Efficacy and Safety Study of Quizartinib (AC220) in Patients \geq 60 Years of Age with FLT3 ITD Positive or Negative Relapsed/Refractory Acute Myeloid Leukemia. *Blood*. 2012;120:48. ([View Abstract](#))

Deol A, Sengsayadeth S, Ahn K, et al. Does FLT3 Mutation Impact Survival After Hematopoietic Stem Cell Transplantation for Acute Myeloid Leukemia: A Center for International Blood and Marrow Transplant Research (CIBMTR) Analysis. *Cancer*. 2016 Jun 17. doi: 10.1002/cncr.30140. [Epub ahead of print] ([View Abstract](#))

Dohner K, Schlenk RF, Habdank M, Scholl C, Rucker FG, Corbacioglu A, et al. Mutant nucleophosmin (NPM1) predicts favorable prognosis in younger adults with acute myeloid leukemia and normal cytogenetics: interaction with other gene mutations. *Blood*. 2005;106(12):3740-6. ([View Abstract](#))

Gale R, Green C, Allen C, et al. The impact of FLT3 internal tandem duplication mutant level, number, size and interaction with NPM1 mutations in a large cohort of young adult patients with acute myeloid leukemia. *Blood* 2008 111:2776-2784 ([View Abstract](#))

Gale RE, Hills R, Kottaridis PD, et al. No evidence that FLT3 status should be considered as an indicator for transplantation in acute myeloid leukemia (AML): an analysis of 1135 patients, excluding acute promyelocytic leukemia, from the UK MRC AML10 and 12 trials. *Blood*. 2005;106(10):3658-3665 ([View Abstract](#))

Kottaridis PD, Gale RE, Linch DC. Flt3 mutations and leukaemia. *Br J Haematol*. 2003;122(4):523-38. ([View Abstract](#))

Kottaridis PD, Gale RE, Frew ME, et al. The presence of a FLT3 internal tandem duplication in patients with acute myeloid leukemia (AML) adds important prognostic information to cytogenetic risk group and response to the first cycle of chemotherapy: analysis of 854 patients from the United Kingdom Medical Research Council AML 10 and 12 trials. *Blood*. 2001 Sep 15;98(6):1752-9. ([View Abstract](#))

Linch DC, Hills RK, Burnett AK, Khwaja A, Gale RE. Impact of FLT3ITD mutant allele level on relapse risk in intermediate-risk acute myeloid leukemia. *Blood*. 2014;124(2):273-6. ([View Abstract](#))

Loghavi S, Zuo Z, Ravandi F, Kantarjian HM, Bueso-Ramos C, Zhang L, et al. Clinical features of de novo acute myeloid leukemia with concurrent DNMT3A, FLT3 and NPM1 mutations. *J Hematol Oncol*. 2014;7:74. ([View Abstract](#))

Ostronoff F, Othus M, Lazenby M, et al. Prognostic Significance of NPM1 Mutations in the Absence of FLT3–Internal Tandem Duplication in Older Patients With Acute Myeloid Leukemia: A SWOG and UK National Cancer Research Institute/Medical Research Council Report. *JCO* Apr 1, 2015;1157-1164. ([View Full Text](#))

Pratz KW, Sato T, Murphy KM, Stine A, Rajkhowa T, Levis M. FLT3-mutant allelic burden and clinical status are predictive of response to FLT3 inhibitors in AML. *Blood*. 2010;115(7):1425-32. ([View Abstract](#))

Ravandi F, Alattar ML, Grunwald MR, et al. Phase II study of azacytidine plus sorafenib in patients with acute myeloid leukemia and FLT-3 internal tandem duplication mutation. *Blood*. 2013;121(23):4655-4662. ([View Abstract](#))

Röllig C, Serve H, Hüttmann A, et al. Addition of sorafenib versus placebo to standard therapy in patients aged 60 years or younger with newly diagnosed acute myeloid leukaemia (SORAML): a multicentre, phase 2, randomised controlled trial. *Lancet Oncol*. 2015 Dec;16(16):1691-9. ([View Abstract](#))

Schnittger S, Bacher U, Kern W, et al. Prognostic impact of FLT3-ITD load in NPM1 mutated acute myeloid leukemia. *Leukemia*. 2011;25, 1297-1304. ([View Abstract](#))

Stone, RM, Mandrekar S, Sanford BL, et al. The multi-kinase inhibitor midostaurin prolongs survival compared with placebo in combination with daunorubicin/cytarabine induction, high-dose cytarabine consolidation, and as maintenance therapy in newly diagnosed acute myeloid leukemia patients age 18-60 with FLT3 mutations: an international prospective randomized placebo-controlled double-blind trial (CALGB 10603/RATIFY). *Blood* 2015 126:6. ([View Abstract](#))

Walter RB, Othus M, Burnett AK, Lowenberg B, Kantarjian HM, Ossenkoppele GJ, et al. Significance of FAB subclassification of "acute myeloid leukemia, NOS" in the 2008 WHO classification: analysis of 5848 newly diagnosed patients. *Blood*. 2013;121(13):2424-31. ([View Abstract](#))